

Characterisation of rock samples collected in the first week of the eruption- trace elements and Pb-isotopes

The trace element concentrations in the lavas varied little throughout the first week of the Geldingadalir eruption, similar to the major elements (Fig. 1, see previous petrology report). The lavas have lower HREE/LREE ratios, and lower abundances of trace elements than historical Reykjanes lavas. This is consistent with the previously reported low TiO₂ content of the lavas.




Fig. 1: Rare earth element concentrations in the Geldingadalir eruption's bulk rock (red) and matrix glass (orange), compared with historical lavas on Reykjanes (light blue). Data is normalised to the primitive mantle composition of Palme & O'Neill (2014) and is shown on a logarithmic scale. The comparison data is taken from Kokfelt et al. (2006), Peate et al. (2009), and Koorneef et al. (2012).

We also report preliminary radiogenic Pb-isotope ratios for material collected on the first day of the eruption (Fig. 2). The Geldingadalir eruption's Pb-isotope ratio values are lower (less radiogenic) than the historical Reykjanes lavas, but sit within the Icelandic array.




Fig. 2: Radiogenic Pb-isotope ratios for glass from Geldingadalir lava sample 20210320-005, collected on the first day of the eruption (orange cross). Historical Reykjanes lavas (Peate et al., 2009) shown as blue circles for comparison. The whole Iceland array is shown in light grey (see references below).

The magmas that are contributing to the Geldingadalir eruption are compositionally distinct from those associated with historical Reykjanes eruptions, in their major element composition (see previous report), their trace element concentrations, and their Pb-isotope



ratios. However, the geochemistry of the lavas is within the range of older magmatism on Reykjanes. Therefore, the shift in the geochemistry identified in lava from the new eruption potentially reflects a new and distinct batch of magma arriving from the mantle beneath Reykjanes.

The Geldingadalir lava trace element concentrations and radiogenic isotopes allow us to identify the mantle sources and mantle melting processes occurring beneath Iceland. The lower concentrations of incompatible trace elements potentially show a larger contribution from higher degree partial mantle melts, sampling more depleted/refractory mantle components. This is further supported by the less-radiogenic Pb-isotope ratios, which indicate a greater contribution from a 'depleted' mantle component, more likely to be sampled by higher degree partial melting.

References

- Peate, D. W., Baker, J. A., Jakobsson, S. P., Waight, T. E., Kent, A. J., Grassineau, N. V., & Skovgaard, A. C. (2009). Historic magmatism on the Reykjanes Peninsula, Iceland: a snap-shot of melt generation at a ridge segment. *Contributions to Mineralogy and Petrology*, 157(3), 359-382.
- Kokfelt, T. F., Hoernle, K. A. J., Hauff, F., Fiebig, J., Werner, R., & Garbe-Schoenberg, D. (2006). Combined trace element and Pb-Nd-Sr-O isotope evidence for recycled oceanic crust (upper and lower) in the Iceland mantle plume. *Journal of Petrology*, 47(9), 1705-1749.
- Koornneef, J. M., Stracke, A., Bourdon, B., Meier, M. A., Jochum, K. P., Stoll, B., & Grönvold, K. (2012). Melting of a two-component source beneath Iceland. *Journal of Petrology*, 53(1), 127-157.
- Palme, H., O'Neill, H. St. C. (2014). Cosmochemical Estimates of Mantle Composition. *Treatise on Geochemistry (Second Edition)*, 3(1), 1-39.

References for Iceland compilation in Fig. 2

- Blichert-Toft, J., Agraniér, A., Andres, M., Kingsley, R., Schilling, J. G., & Albarède, F. (2005). Geochemical segmentation of the Mid-Atlantic Ridge north of Iceland and ridge-hot spot interaction in the North Atlantic. *Geochemistry, Geophysics, Geosystems*, 6(1).
- Chauvel, C., & Hémond, C. (2000). Melting of a complete section of recycled oceanic crust: Trace element and Pb isotopic evidence from Iceland. *Geochemistry, Geophysics, Geosystems*, 1(2).
- Elliott, T. R., Hawkesworth, C. J., & Grönvold, K. (1991). Dynamic melting of the Iceland plume. *Nature*, 351(6323), 201-206.
- Furman, T., Frey, F. A., & Park, K. H. (1991). Chemical constraints on the petrogenesis of mildly alkaline lavas from Vestmannaeyjar, Iceland: the Eldfell (1973) and Surtsey (1963-1967) eruptions. *Contributions to Mineralogy and Petrology*, 109(1), 19-37.
- Halldorsson, S. A., Oskarsson, N., Grönvold, K., Sigurdsson, G., Sverrisdóttir, G., & Steinthorsson, S. (2008). Isotopic-heterogeneity of the Thjorsa lava—implications for mantle sources and crustal processes within the Eastern Rift Zone, Iceland. *Chemical Geology*, 255(3-4), 305-316.
- Hards, V. L., Kempton, P. D., & Thompson, R. N. (1995). The heterogeneous Iceland plume: new insights from the alkaline basalts of the Snaefell volcanic centre. *Journal of the Geological Society*, 152(6), 1003-1009.
- Hards, V. L., Kempton, P. D., Thompson, R. N., & Greenwood, P. B. (2000). The magmatic evolution of the Snaefell volcanic centre; an example of volcanism during incipient rifting in Iceland. *Journal of Volcanology and Geothermal Research*, 99(1-4), 97-121.
- Kempton, P. D., Fitton, J. G., Saunders, A. D., Nowell, G. M., Taylor, R. N., Hardarson, B. S., & Pearson, G. (2000). The Iceland plume in space and time: a Sr-Nd-Pb-Hf study of the North Atlantic rifted margin. *Earth and Planetary Science Letters*, 177(3-4), 255-271.
- Kitagawa, H., Kobayashi, K., Makishima, A., & Nakamura, E. (2008). Multiple pulses of the mantle plume: evidence from Tertiary Icelandic lavas. *Journal of Petrology*, 49(7), 1365-1396.
- Kokfelt, T. F., Hoernle, K. A. J., Hauff, F., Fiebig, J., Werner, R., & Garbe-Schoenberg, D. (2006). Combined trace element and Pb-Nd-Sr-O isotope evidence for recycled oceanic crust (upper and lower) in the Iceland mantle plume. *Journal of Petrology*, 47(9), 1705-1749.
- Kuritani, T., Yokoyama, T., Kitagawa, H., Kobayashi, K., & Nakamura, E. (2011). Geochemical evolution of historical lavas from Askja Volcano, Iceland: implications for mechanisms and timescales of magmatic differentiation. *Geochimica et Cosmochimica Acta*, 75(2), 570-587.
- Peate, D. W., Baker, J. A., Jakobsson, S. P., Waight, T. E., Kent, A. J., Grassineau, N. V., & Skovgaard, A. C. (2009). Historic magmatism on the Reykjanes Peninsula, Iceland: a snap-shot of melt generation at a ridge segment. *Contributions to Mineralogy and Petrology*, 157(3), 359-382.
- Peate, D. W., Breddam, K., Baker, J. A., Kurz, M. D., Barker, A. K., Prestvik, T., ... & Skovgaard, A. C. (2010). Compositional characteristics and spatial distribution of enriched Icelandic mantle components. *Journal of Petrology*, 51(7), 1447-1475.
- Prestvik, T., Goldberg, S., Karlsson, H., & Grönvold, K. (2001). Anomalous strontium and lead isotope signatures in the off-rift Öraefajökull central volcano in south-east Iceland: Evidence for enriched endmember (s) of the Iceland mantle plume?. *Earth and Planetary Science Letters*, 190(3-4), 211-220.
- Schilling, J. G., Kingsley, R., Fontignie, D., Poreda, R., & Xue, S. (1999). Dispersion of the Jan Mayen and Iceland mantle plumes in the Arctic: A He-Pb-Nd-Sr isotope tracer study of basalts from the Kolbeinsey, Mohns, and Knipovich Ridges. *Journal of Geophysical Research: Solid Earth*, 104(B5), 10543-10569.
- Shorttle, O., MacLennan, J., & Piotrowski, A. M. (2013). Geochemical provincialism in the Iceland plume. *Geochimica et Cosmochimica Acta*, 122, 363-397.
- Sims, K. W., MacLennan, J., Blichert-Toft, J., Mervine, E. M., Blusztajn, J., & Grönvold, K. (2013). Short length scale mantle heterogeneity beneath Iceland probed by glacial modulation of melting. *Earth and Planetary Science Letters*, 379, 146-157.
- Stracke, A., Zindler, A., Salters, V. J., McKenzie, D., & Grönvold, K. (2003). The dynamics of melting beneath Theistareykir, northern Iceland. *Geochemistry, Geophysics, Geosystems*, 4(10).
- Thirlwall, M. F., Gee, M. A. M., Taylor, R. N., & Murton, B. J. (2004). Mantle components in Iceland and adjacent ridges investigated using double-spike Pb isotope ratios. *Geochimica et Cosmochimica Acta*, 68(2), 361-386.